LLE Review Quarterly Report



Contents

Director's Remarks	iii
IN BRIEF	vii
Inertial Confinement Fusion	
Diagnosing Low-Mode ($\ell < 6$) and Mid-Mode ($6 \le \ell \le 60$) Asymmetries in the Post-Stagnation Phase of Cryogenic Implosions on OMEGA	1
Effect of Overlapping Laser Beams and Density Scale Length in Laser–Plasma Instability Experiments on OMEGA EP	4
Optimization of Irradiation Configuration Using Spherical <i>t</i> Designs for Laser-Direct-Drive Interial Confinement Fusion	7
Validation of Ray-Based Cross-Beam Energy Transfer Models	10
Diagnosis of the Imploding Shell Asymmetry in Polar-Direct-Drive Deuterium–Tritium Cryogenic Target Implosions on OMEGA	13
PLASMA AND ULTRAFAST PHYSICS	
Quantitative Proton Radiography and Shadowgraphy for Arbitrary Intensities	18
Direct Measurement of the Return-Current Instability	22
Analysis Methods for Electron Radiography Based on Laser-Plasma Accelerators	25
High-Energy-Density Physics	
A First-Principles Study of L-Shell Iron and Chromium Opacity at Stellar Interior Temperatures	28

DIAGNOSTIC SCIENCE AND DETECTORS

High-Resolution X-Ray Spectrometer for X-Ray Absorption Fine Structure Spectroscopy	31
(Cd,Mg)Te Crystals for Picosecond-Response Optical-to-X-Ray Radiation Detectors	34
Exact Solutions for the Electromagnetic Fields of a Flying Focus	38
LASER TECHNOLOGY AND DEVELOPMENT	
Methods and Apparatus for Comprehensive Characterization of Performance Attributes and Limits of Ultrafast Laser Optics	41
Design and Demonstration of the "Flow-Cell" Integrated-Flow, Active-Cooling Substrate for High-Average-Power Substrates	44
Free-Standing Thin-Membrane, Zero B-Integral Beam Splitters	48
EDUCATION AND OUTREACH	
Petawatt Laser Systems	51
LLE's Summer High School Research Program	54
2022 BEST Student and Teacher Research Program	57
LASER FACILITY	
FY22 Q4 Laser Facility Report	62
Groundbreaking for the LLE Building Expansion	64
ANNUAL REPORTS	
FY22 Laser Facility Report	67
Education Summary	71
External Users' Programs	79
FY22 National Laser Users' Facility Program	83
FY22 Laboratory Basic Science Program	126
FY22 LaserNetUS Program	156
PUBLICATIONS AND CONFERENCE PRESENTATIONS	162

Director's Remarks

C. Deeney

Director, Laboratory for Laser Energetics

The Laboratory for Laser Energetics (LLE) is funded primarily by the U.S. Department of Energy (DOE) National Nuclear Security Administration's (NNSA's) Office of Experimental Sciences Inertial Confinement Fusion (ICF) Program through a five-year Cooperative Agreement. The fiscal year ending September 2022 (FY22) comprised the fourth year of work under DOE/NNSA Cooperative Agreement No. DE-NA0003856. The Laboratory is also sponsored by the New York State Energy Research Development Authority and other federal agencies such as the DOE Office of Science and the National Science Foundation. I was honored to become the fifth Director of LLE, succeeding Dr. E. Michael Campbell who retired in March 2022.

The year 2022 was exciting for the National ICF Program and LLE. Ignition was achieved at the National Ignition Facility for the first time, producing more energy out of the implosion than the input laser energy to the target. LLE has made many contributions to the pursuit of ignition, including laser technologies, diagnostics, simulation capabilities, physics understanding, and student education. Groundbreaking for the construction of a new \$42M addition to the LLE Complex (funded by the University of Rochester) occurred in August 2022. LLE submitted a proposal to renew the Cooperative Agreement with DOE/NNSA for FY24–FY28.

This report summarizes work conducted at LLE during FY22 that includes research on the ICF and High-Energy-Density (HED) science campaigns; laser, optical materials and advanced technology development; operation of the Omega Laser Facility for the ICF and HED campaigns, the National Laser Users' Facility, Laboratory Basic Science Program, and other external users, including the newly established LaserNetUS supported by the DOE Office of Fusion Energy Sciences; and programs focusing on the education of high school, undergraduate, and graduate students.

- The Omega Laser Facility conducted 2110 target shots in FY22, close to the average annual number of shots since FY14 with nearly 60% of the shots performed for external users. These experiments are critical to LLE and the national community achieving progress in the ICF/HED Physics mission, in advancing science, and in training students.
- Mission Impact: All of the completion criteria for the 15 NNSA Level II milestones were achieved.
- Scientific Output: One hundred and one technical manuscripts were published in peer-reviewed scientific journals during FY22.
- Sixteen Ph.D. degrees were conferred on graduate students during 2022 whose primary research was performed at the Laboratory either as students of the University of Rochester or as students of other collaborating user institutions who performed experiments or worked with scientists at the Omega Laser Facility.
- One University of Rochester Ph.D. graduate (Alison Christopherson) won the American Physical Society Division of Plasma Physics Marshall N. Rosenbluth Outstanding Doctoral Thesis Award.

- Radha Bahukutumbi, a Distinguished Scientist in the Theory Division, received the Leadership Award at the annual meeting of the Fusion Power Associates.
- Steve Ivancic, head of the Diagnostic Development and Integration Group, received a Defense Programs Award of Excellence from Lawrence Livermore National Laboratory.

Key technical highlights included:

- A series of high in-flight aspect ratio and high-velocity implosions on OMEGA produced yields between 2.6×10^{14} and 3.1×10^{14} , all above the previous yield record. The highest yield implosion, when corrected for ³He buildup using statistical modeling, would have produced a yield of nearly 1 kJ, roughly the same as the shell kinetic energy. These direct-drive implosions enable Ph.D. research on ICF and open opportunities for future ICF optimization.
- Layered DT implosions with silicon dopants performed significantly better (higher yields) than comparable targets without dopants, confirming that the dopant suppresses imprint while only modestly increasing the shell adiabat via x-ray preheat.
- The Lawrence Livermore National Laboratory (LLNL)/LLE Neutron Sources collaboration conducted a polar-direct-drive exploding-pusher experiment at the National Ignition Facility that produced a record laser-direct-drive (LDD) yield of 1.6×10^{16} with a laser pulse of 1.6 MJ.
- The proof-of-concept for the LLE dynamic-shell ignition design, where pulse shaping dynamically creates a high-density shell with a DT wetted-foam target, was demonstrated using 3-D-printed foam capsules.
- Constrained by integrated implosion measurements, spectroscopic observables revealed that a density functional theory (DFT)based kinetic model developed by LLE scientists (*VERITAS*) reproduces a majority of the emission and absorption features observed in experiments, while the traditional collisional-radiative-equilibrium treatment with *ad hoc* continuum lowering does not, suggesting that self-consistent treatment of dense plasma effects is needed at very high pressures.
- In collaboration with LLNL, LLE scientists showed that diamond can precipitate from twice-shocked CH polymers (e.g., ICF ablators) in the presence of oxygen, nitrogen, and chlorine.
- An LLE–LLNL team conclusively demonstrated species separation in the release of strongly shocked CH, finding that the hydrogen ions carry 10 to 100× more mass than the carbon ions to large distances, confirming a hypothesis developed to understand the seeds of instability growth in plastic ICF ablators.
- Based on x-ray diffraction measurements of sevenfold compressed sodium, a team from LLE and LLNL observed the hP4 electride phase at a pressure of 480 GPa and a temperature of 3000 K (a regime where core electron overlap was thought to stabilize against this phase), indicating that electride formation is possible on nanosecond time scales at high temperatures.
- A decades-old discrepancy between theory and measurement on the shock-induced metallization of polystyrene was resolved using a new exchange correlation functional that properly captures the shock-induced dissociation of carbon and correctly predicts the band-gap closing behavior at 1 to 2 Mbar.
- New DFT calculations of Fe and Cr opacity at stellar interior densities and temperatures did not resolve the iron opacity "mystery" identified some time ago at Sandia National Laboratories' Z Pulsed Power Facility. The new calculations showed good agreement with the measured Cr opacity but did not agree with Z's measurements of the Fe opacity.

- Recent experiments showed that micron-scale x-ray radiography is possible for a wide range of flow visualizations at ultrahigh pressures using a new LLE-developed Fresnel zone plate imaging diagnostic.
- The first electron radiographs of a test object were acquired (in collaboration with Los Alamos National Laboratory) using a laser-plasma accelerator platform on OMEGA EP.
- The 13th Omega Laser Facility Users Group (OLUG) Workshop, held 27–29 April 2022 at LLE, attracted nearly 200 researchers from around the world.
- The 24th Topical Conference on High-Temperature Plasma Diagnostics (HTPD), hosted by LLE in Rochester, NY, attracted 370 registered participants. The HTPD conference showcased the latest in advanced instrumentation in the fields of magnetic confinement fusion, inertial confinement fusion, high-energy-density plasmas, space plasmas, astrophysics, and industrial applications.
- The 50th Anomalous Absorption Conference 2022 hosted and organized by LLE was held 5–10 June in Skytop, PA. Conference topics included theoretical and experimental studies in parametric instabilities, radiation hydrodynamics, particle acceleration, HED physics, short-pulse laser-matter interactions, and ICF.
- A classified report on the scope of applications within the Stockpile Stewardship Program for platforms that rely on LDD was submitted to the NNSA Office of Experimental Science, completing one of the milestones in the ICF five-year plan.

As the fifth Director working with the senior leadership team, we have set our vision to be "the leading academic institution advancing laser technologies, inertial confinement fusion, and high-energy-density science at scale." I hope you agree that FY22 met this vision.

In Brief

This volume of LLE Review 172 covers the period from July–September 2022. Articles appearing in this volume are the principal summarized results for long-form research articles. Readers seeking a more-detailed account of research activities are invited to seek out the primary materials appearing in print, detailed in the publications and presentations section at the end of this volume.

Highlights of research presented in this volume include:

- J. Baltazar *et al.* develop a technique to diagnose the low- and mid-mode asymmetries in the post-stagnation phase of a deuterium–tritium (DT) cryogenic implosion with the aim of relating the post-stagnation phase measurement to the shell breakup caused by an increase in beam overlap perturbations due to a lower R_b/R_t (p. 1).
- M. J. Rosenberg *et al.* analyze laser–plasma instability mitigation techniques for direct-drive inertial confinement fusion (ICF) by examining stimulated Raman scattering and two-plasmon–decay instabilities, assessing their dependence on density scale length and overlapping laser beam geometry critical to ICF ignition/fusion gain (p. 4).
- A. Shvydky *et al.* propose a new approach in producing uniform beam intensity distributions applied on a sphere for laserdirect-drive ICF system configurations based on mathematical spherical *t* designs to find optimal beam configurations, targeting a reduction of spherical harmonics and nonuniformity modes (p. 7).
- R. K. Follett *et al.* develop a series of test cases for validation of ray-based cross-beam energy transfer (CBET) models against wave-based calculations. Comparisons between various ray-based models show that an etalon integral field reconstruction with a coherent caustic correction and caustic gain truncation is the preferred ray-based CBET model, with equal to improved performance without increasing computational cost (p. 10).
- T. R. Joshi *et al.* extend the application of the x-ray self-emission imaging technique to polar-direct-drive ICF implosions with cryogenically layered DT targets on OMEGA, diagnosing in-flight asymmetries (p. 13).
- J. R. Davies, P. V. Heuer, and A. F. A. Bott develop an electrostatic particle-in-cell (PIC) algorithm that was compared to algorithms for charged-particle radiography and shadowgraphy used in laser-plasma experiments, deducing a Monge–Ampère code applied as first order, and a PIC code if the Monge–Ampère code fails or requires a subsequent, more-accurate inversion (p. 18).
- A. L. Milder *et al.* present first measurements of the threshold and linear growth rate of the return-current instability driven by electron heat flux, where characterization of the plasma conditions show that the return-current instability occurs concurrently with nonlocal transport (p. 22).
- G. Bruhaug *et al.* present a method to determine the resolution for laser-plasma acceleration electron radiography that include accounting for drive laser image distortions, quantifying plasma-generated electric fields in laser-ablated targets by measuring electron radiography image feature sizes, and estimating field strength needed to produce those features (p. 25).
- V. V. Karasiev *et al.* apply first-principles density functional theory to calculate optical properties (mass-absorption coefficient and opacity) of Cr and Fe at stellar interior temperatures to explore whether *ab initio* calculations can resolve the disagreement between previous atomic physics calculations and measured data (p. 28).
- D. A. Chin *et al.* designed and built two extended x-ray absorption fine structure (EXAFS) flat crystal x-ray spectrometers (EFX's) for high-resolution x-ray spectroscopy over a large energy range with flexible, on-shot energy dispersion calibration capabilities, enabling x-ray absorption near-edge spectroscopy measurements to be made on OMEGA (p. 31).

- J. Cheng *et al.* demonstrate a promising candidate for ultrafast optical/near-infrared to x-ray radiation detector applications with a photodetector sensitive to both optical and x-ray picosecond pulses based on in-house grown cadmium magnesium telluride (Cd,Mg) Te single crystal (p. 34).
- D. Ramsey *et al.* derive exact solutions to Maxwell's equations for the electromagnetic fields of a constant-velocity flying-focus pulse and identified small differences to the paraxial solutions for a wide range of parameters, justifying paraxial solutions for many applications in many regimes, such as a spatiotemporal pulse-shaping technique (p. 38).
- K. R. P. Kafka, T. Z. Kosc, and S. G. Demos assess operational performance limits of optics and aspects required for the development of improved optic designs/reliable operational fluence limits for ultrashort optical performance characterization beyond the standard laser-induced-damage threshold testing system specifications that only characterize the damage-initiation threshold (p. 41).
- E. P. Power *et al.* design, simulate, fabricate, and test a prototype flow-cell, integrated-cooling substrate built using cordierite ceramic and demonstrate average power handling up to 3.88-W/cm² absorbed power density with 54-nm peak-to-valley deformation in a sub-aperture test, with a <30 s observed mechanical stabilization time scale (p. 44).
- M. Romo-Gonzalez and R. Boni test a zero *B*-integral pellicle beam splitter composed of uncoated nitrocellulose, having a maximum reflectivity of ~30% with negligible absorption in the near-infrared (p. 48).
- L. J. Waxer, J. Bromage, and B. E. Kruschwitz provide an overview of the various topics they wrote in the SPIE Spotlight series e-book on petawatt laser systems, introducing the reader to the laser science and technology underpinning petawatt laser systems and technological advances required for state-of-the-art high-intensity laser performance (p. 51).
- R. S. Craxton summarizes the 33rd LLE Summer High School Research Program (p. 54). Sixteen students were invited from Rochester-area high schools to participate in the lab's state-of-the art research environment.
- T. J. Kessler, M. Romo-Gonzalez, and R. Ghosh report on the Broad Exposure to Science and Technology (BEST) Program, a research program designed to engage teachers and students from historically marginalized experiences in various aspects of science and technology that support LLE's laser science and applications research (p. 57).
- J. Puth et al. summarize operations of the Omega Laser Facility during the fourth quarter of FY22 (p. 62).
- M. J. Shoup III highlights the groundbreaking and early days of the LLE building expansion (p. 64).
- This volume concludes with the FY22 Laser Facility Report, the FY22 Education Summary, and overviews of the National Laser Users' Facility, Laboratory Basic Science, and LaserNetUS Programs.

Nickolaos Savidis Editor